

# Factors affecting the size of a mosquito population in a favourable environment

**P. Cailly<sup>1</sup>, A. Tran<sup>2</sup>, T. Balenghien<sup>3</sup>, C. Toty<sup>4,5</sup>, P. Ezanno<sup>1</sup>**

<sup>1</sup>INRA, Oniris, UMR1300 Bio-agression, Epidémiologie et Analyse de Risques,

<sup>2</sup>CIRAD, UPR AGIRs Animal et Gestion Intégrée des Risques,

<sup>3</sup>CIRAD, UMR Contrôle des maladies,



<sup>4</sup>IRD, UR016 Caractérisation et Contrôle des Populations de Vecteurs,

<sup>5</sup>Centre de Recherche et de Veille sur les maladies émergentes dans l'Océan Indien



# Why mosquitoes?

- Pathogenic agents of major vector born diseases
  - viruses (e.g. West Nile virus), parasites (e.g. *Plasmodium falciparum*)
  - largely distributed across the world
  - public health & animal health
  - health, ecological, socioeconomic & political consequences



➔ active vectors necessary for the epidemiological cycle of these diseases

# Why mosquitoes?

---

- Efficient anti-vector fight strategies require to well know:
  - mosquito life cycle & behaviour
  - related to their biotope (breeding site, ...)
  - related to the variation of climatic factors (temperature, humidity, ...)
- Mosquitoes brave these strategies up to now: malaria and dengue do not weaken
- An integrative approach by modelling proves to be necessary for better:
  - understanding the dynamics of a mosquito vector population
  - describing the mosquito abundance
  - identifying the most influential parameters = potential control points of the population




# Objective

---

Identifying by modelling the factors affecting the size of a mosquito population located in a favourable environment

# Modelling approach

- Mechanistic model
- Adapted to *Anopheles* of Camargue region, France
- Driven by climate
- Lasting several years





# Modelling approach

- Nulliparous into diapause during winter
- Development parameters & mortality rates depend on temperature
- Model output

- General output:


Dynamic abundance in adults over time

- Aggregated output:



# Sensitivity analysis of the model

- Quantify the influence of input values on the outputs




- describe the model parameters for which the model is the most sensitive
  - ➔ Influent parameters are potential control points of the biological system
- Confront different methods to identify influential parameters in a robust way



# Results

- Model confrontation to independent field data
  - Number of host-seeking females




→ Our model predicts correctly a mean dynamics of mosquito populations



# Results


## Sensitivity analysis

- No interaction
- Influent parameters:



# Scenario of control strategies

- Bio-larvicide, similar to Bti, persistent during 7 days
- Different proportions of the sprayed surface & 2 strategies tested
  - Sprayed at regular time intervals
  - Sprayed when mosquito abundance exceeds a threshold value



## Discussion & perspectives

- Contribution of our model of mosquito population dynamics
  - Knowing the influential parameters of the model orientates future research efforts on the control of mosquitoes
  - The model is a tool to test control strategies
- Valid results for a homogeneous environment
  - Not restricted in hosts, rest shelters, breeding sites
- Environmental factors can be heterogeneous
  - ➔ Population can be spatially structured
- In these conditions:
  - ➔ Spatial model is the next step




# Thank you for your attention ...

---



**Questions?**





● Scenario corresponding to sprayed surface=0.45 , number of treatment=5

